

Angelini Pharma Procedure for Use - Antisepsis Preparation for Graft/AVF Cannulation

SUPPLIES

ExSept Plus Solution
Two Alcohol Prep Pads
Two 2x2 Gauze Pads
Chux Sheet
Non-Sterile Gloves
Face Mask



PROCEDURE STEPS

Have the patient wash their access site with soap and water prior to sitting in the chair.

Staff members don PPE and hand wash.

On a chux sheet, carefully open two packages of 2x2 gauze pads, keeping pads in their sterile wrappers. Saturate both sets of 2x2 gauze pads with ample amounts of ExSept Plus (3-5ml).

Don clean, non-sterile gloves.

Take one saturated ExSept Plus 2x2 gauze pad and clean the cannulation site starting at the center and moving in a continuous circle outward to a circumference of 2 inches.

Repeat this step with the second cannulation site.

Allow skin to air dry.

Open alcohol prep pad and clean cannulation site for 15 seconds, using a scrubbing motion.

Repeat this step with the second cannulation site using a new alcohol prep pads.

Allow to air dry.

Initiate dialysis treatment per facility protocol.



CONTACT US

Angelini Pharma Inc.
8322 Helgerman Court - Gaithersburg - MD 20877 -USA
1 (800) 726-2308
www.Angelini-us.com

10-015-00 Rev 00 10/14